
How To Make Yourself Poop

*How To
Make
Yourself
Poop*

*Downloaded from
www.everybodystreet.com
by Breanna & Michaela*

UNDERSTANDING THE DISCOMFORT OF CONSTIPATION

Feeling backed up, bloated, and uncomfortable is a common experience that can disrupt your entire day. The sensation of needing to go but being unable to is frustrating, and often, people find themselves urgently searching for ways to get things moving. If you are looking for effective and safe methods on how to make yourself poop, you are not alone.

Constipation affects people of all ages and backgrounds, but the good news is that in most cases, it can be relieved with simple, natural adjustments. This comprehensive guide will walk you through evidence-based strategies to relieve constipation and get your digestive system back on track, focusing on long-term digestive health rather than just a quick fix.

Before diving into solutions, it helps to understand what constipation actually is. Medically, it is often defined as having fewer than three bowel

movements per week. However, constipation also involves symptoms like passing hard, dry stools, straining during bowel movements, or feeling like you haven't completely emptied your bowels. The causes can range from diet and hydration to stress, lack of physical activity, or even ignoring the urge to go. By addressing these root causes, you can find lasting relief. This article will explore a step-by-step approach to how to make yourself poop, emphasizing the most effective and gentle methods available.

IMMEDIATE DIETARY CHANGES TO STIMULATE A BOWEL MOVEMENT

Increase Your Fiber Intake Strategically

Fiber is the single most important nutrient for healthy bowel movements. It works by adding bulk to your stool and softening it, which makes it easier to pass. There are two types of fiber, and both play a role in how to make yourself poop.

- **Soluble Fiber:**
This type dissolves in water to form a gel-like substance. It

helps soften stool. Good sources include oats, barley, nuts, seeds, beans, lentils, and fruits like apples and carrots. Psyllium husk, the main ingredient in many fiber supplements like Metamucil, is a powerful soluble fiber.

- **Insoluble Fiber:**

This type does not dissolve in water. It adds bulk to the stool and acts like a broom, helping food pass more quickly through your stomach and intestines. Good sources include whole-wheat flour, wheat bran, nuts, beans, and

vegetables like cauliflower, green beans, and potatoes (with skin).

Important Note: If you are not used to a high-fiber diet, increase your intake gradually over a few days. Adding too much fiber too quickly can cause gas, bloating, and cramping, which will make the problem worse. Always drink plenty of water when increasing fiber, as fiber works by absorbing water. Without enough water, fiber can actually worsen constipation.

Hydrate, Hydrate,

Hydrate

Water is essential for healthy digestion. Your colon pulls water from your body to soften stool. When you are dehydrated, the colon absorbs too much water, leading to hard, dry stools that are difficult to pass. Aim for at least eight 8-ounce glasses of water per day, and more if you are active or live in a hot climate. Warm liquids, in particular, can be very effective. Drinking a warm glass of water first thing in the morning or sipping hot herbal tea can stimulate the digestive tract and trigger a bowel movement. Many people find that warm water with lemon is an excellent morning habit for regularity.

Incorporate Natural Laxative Foods

Certain foods are particularly well-known for their ability to get things moving. Adding these to your diet can provide gentle relief when you are struggling with how to make yourself poop.

- **Prunes (Dried Plums) and Prune Juice:**

This is the classic natural remedy for a reason.

Prunes are rich in fiber, particularly insoluble fiber, and also contain sorbitol, a sugar alcohol that acts as a natural

laxative by drawing water into the intestines. Eating three to four prunes or drinking a small glass of prune juice can often stimulate a bowel movement within a few hours.

- **Kiwi Fruit:**

Studies have shown that eating two kiwifruits per day can increase stool frequency and soften stool consistency. It contains a unique enzyme called actinidin that helps with digestion, as well as fiber and other compounds that promote regularity.

- **Apples and**

Pears: These fruits are high in pectin, a type of soluble fiber that works similarly to psyllium. They also contain natural sugars and water that can help soften stool. Eat them with the skin on for maximum fiber benefit.

- **Leafy Greens:**

Spinach, kale, collard greens, and Swiss chard are excellent sources of insoluble fiber and also contain magnesium, which can help relax the bowel muscles.

- **Chia Seeds and Flaxseeds:**

These tiny seeds are packed with soluble fiber. When mixed with

liquid, they form a gel that can help soften stool. Add a tablespoon of ground flaxseed or whole chia seeds to a smoothie, oatmeal, or yogurt. Remember to drink extra water when consuming seeds.

- **Coffee (Especially Caffeinated):**

For many people, a morning cup of coffee stimulates the colon. The caffeine in coffee can trigger the release of hormones that increase muscular contractions in the gut, urging a bowel movement. Just be mindful that

too much coffee can lead to dehydration for some people, so balance it with water.

LIFESTYLE ADJUSTMENTS FOR LONG-TERM REGULARITY

Establish a Regular Bathroom Routine

Your body thrives on routine. Your digestive system is no different. Try to set aside a specific time each day for a bowel movement. For most people, the morning after breakfast is the most natural time, as eating stimulates the

gastrocolic reflex - a signal from your stomach to your colon to move waste along. Even if you do not feel the urge, sit on the toilet for 5 to 10 minutes at the same time each day. Over time, your body will learn to follow the routine. Do not ignore the urge to go when it strikes. Holding it in can lead to the stool becoming harder and more difficult to pass.

Physical Activity and Movemen t

Exercise is not just for your heart and muscles; it is fantastic

for your digestion as well. Physical activity increases blood flow to your digestive organs and stimulates the natural contractions of your intestines (peristalsis). Even moderate exercise like a 20-minute walk after a meal can make a significant difference in how to make yourself poop. Yoga in particular can be helpful. Twisting poses, such as a seated spinal twist, and poses that compress the abdomen, like child's pose, are thought to massage the colon and stimulate bowel activity. Aim for at least 30 minutes of moderate physical activity most days of the week.

Manage Stress Levels

The gut-brain connection is powerful. Stress, anxiety, and tension can wreak havoc on your digestive system. Your gut is often called your "second brain" because it is lined with millions of nerve cells that communicate directly with your brain. When you are stressed, your body's fight-or-flight response can slow down or stop digestion, leading to constipation. Practicing stress-reduction techniques like deep breathing, meditation, yoga, or simply taking quiet time for yourself can help regulate your digestion. Even five

minutes of deep, diaphragmatic breathing (belly breathing) can help calm your nervous system and encourage a bowel movement.

GENTLE OVER-THE-COUNTER AND NATURAL REMEDIES

If dietary and lifestyle changes are not enough, there are several safe, over-the-counter options to consider. Always talk to your doctor or a pharmacist before starting any new supplement or medication, especially if you have a medical condition or take other medications.

Osmotic

Laxatives

Osmotic laxatives work by drawing water into the colon from surrounding tissues, which softens the stool and stimulates a bowel movement. They are generally considered gentle and safe for short-term use. Common examples include:

- **Polyethylene Glycol (PEG) - MiraLax:** This is a very popular and well-tolerated option. It is tasteless, odorless, and dissolves easily in water. It usually works within 24 to 48 hours.
- **Magnesium-Based Products:** Milk of Magnesia

(magnesium hydroxide) and magnesium citrate are other osmotic laxatives. Magnesium also has a muscle-relaxing effect, which can help relax the bowel wall. They often work more quickly than PEG, sometimes within 30 minutes to 6 hours.

- **Lactulose:** This is a synthetic sugar that works as an osmotic laxative. It is often prescribed for chronic constipation but is also available over the counter in some places.

Stimulant Laxatives

Stool Softeners

Stimulant laxatives work by directly stimulating the nerves in the colon wall to contract, pushing the stool through. They are more powerful than osmotic laxatives and can cause cramping. They are not recommended for long-term use as they can lead to dependency and disrupt the natural function of your colon. Examples include senna (Senokot) and bisacodyl (Dulcolax). They often work within 6 to 12 hours and are best used for occasional, short-term relief.

A stool softener, such as docusate sodium (Colace), is not a laxative in the sense that it does not stimulate a bowel movement. Instead, it adds moisture to the stool, making it softer and easier to pass. It is most helpful for people who are straining due to hard stools, such as after surgery or childbirth. It usually takes 1 to 3 days to work.

Fiber Supplements

As mentioned earlier, fiber supplements like

psyllium (Metamucil), methylcellulose (Citrucel), or calcium polycarbophil (FiberCon) can be very effective for long-term management. They work gently by adding bulk and softening the stool. It is crucial to drink plenty of water when taking these. They can take 12 to 72 hours to produce a bowel movement.

NATURAL AND HOME REMEDIES WORTH TRYING

Before reaching for medication, many people find relief with simple home remedies that are backed by tradition and often by science.

- **Warm Water and Lemon:** A warm glass of water with fresh lemon juice first

thing in the morning can stimulate digestion and hydrate the colon.

- **Castor Oil:** This is a powerful natural laxative. Taking a small amount (a teaspoon) can produce a bowel movement within a few hours. However, it has a strong taste and can cause cramps. Use it only occasionally and with caution.

- **Aloe Vera Juice:** Aloe vera latex, found just under the leaf skin, contains compounds called anthraquinones that have a laxative effect. However,

commercially available aloe juice often has these compounds removed. If you do find aloe vera juice for digestive health, start with a small amount. It can cause cramping and electrolyte imbalances if overused.

- **Olive Oil or Flaxseed Oil:** A tablespoon of olive oil or flaxseed oil on an empty stomach or in a meal can help lubricate the digestive tract and soften stool. It is a mild and safe option.

- **Probiotics:** These are beneficial bacteria that live in your gut. An

imbalance in gut bacteria can contribute to constipation. Eating fermented foods like yogurt, kefir, sauerkraut, kimchi, and miso, or taking a high-quality probiotic supplement, can help restore balance and improve regularity over time.

A simple, step-by-step morning routine for constipation relief could look like this: Wake up, drink a warm glass of water with lemon. Go for a 10-minute walk or do some gentle stretching. Eat a breakfast rich in fiber, such as oatmeal with berries and a tablespoon of ground flaxseed, plus a whole piece of fruit. After

breakfast, sit on the toilet for 5 to 10 minutes, even if you do not feel the urge. This routine helps set your body's clock for a predictable bowel movement.

WHEN TO SEEK PROFESSIONAL MEDICAL ADVICE

While occasional

constipation is normal and usually easy to treat, there are times when you should see a doctor. Do not rely solely on self-treatment if you experience any of the following:

- Constipation that lasts for more than three