
Review Sheet Exercise 9 The Appendicular Skeleton Answers

*Review Sheet Exercise 9
The Appendicular
Skeleton Answers*

*Downloaded from
www.everybodystreet.com
by Dyer & Cameron*

MASTERING THE APPENDICULAR SKELETON: A COMPREHENSIVE GUIDE TO REVIEW SHEET EXERCISE 9

For students diving into human anatomy and physiology, few tasks are as foundational as understanding the skeletal system. While the axial skeleton provides the central framework, the

appendicular skeleton is what allows us to interact with the world. It is the system of bones that facilitates movement, manipulation, and locomotion. If you are working through a standard anatomy lab manual, you have likely encountered **Review Sheet Exercise 9 The Appendicular Skeleton Answers**. This article serves as a deep dive into that very topic, providing a clear, structured overview of the bones, landmarks, and concepts you

need to master. Whether you are preparing for a lab practical or simply seeking a clearer understanding, this guide will help you connect the dots between the bones of your limbs and girdles.

The appendicular skeleton is composed of 126 bones, divided into two major groups: the pectoral (shoulder) girdle and the upper limbs, and the pelvic (hip) girdle and the lower limbs. Understanding these divisions is the first step. Review Sheet Exercise 9 typically asks students to identify specific bones, label anatomical features, and explain how these structures function together. Let us break down each section systematically, ensuring you have the knowledge to confidently fill in those answers.

THE PECTORAL GIRDLE: CONNECTING THE UPPER LIMBS

The pectoral girdle is the bony ring that attaches the upper limbs to the axial skeleton. It is unique because it provides a great deal of mobility at the cost of stability. This girdle consists of only two bones on each side: the clavicle and the scapula.

The Clavicle: The Collarbone

The clavicle is an S-shaped, long bone that serves as a strut between the sternum and the scapula. It is the only direct bony attachment of the upper limb to the trunk. When studying

Review Sheet Exercise 9 The Appendicular Skeleton Answers, you will often be asked to identify the sternal end (which is rounded and articulates with the manubrium) and the acromial end (which is flatter and articulates with the scapula). This bone is frequently fractured due to its superficial position and the force transmitted through it during a fall.

The Scapula: The Shoulder Blade

The scapula is a flat, triangular bone located on the posterior aspect of the rib cage. It is not directly attached to the axial skeleton by a joint; it is held in place by muscles. Key landmarks you

should be able to locate include:

- **The spine:** A prominent ridge running diagonally across the posterior surface.
- **The acromion:** The lateral extension of the spine that forms the highest point of the shoulder and articulates with the clavicle.
- **The coracoid process:** A beak-like projection on the anterior surface, serving as an attachment point for muscles.
- **The glenoid cavity:** A shallow, socket-like structure that articulates with the head of the humerus to form the shoulder joint.
- **The supraspinous and infraspinous fossae:**

Depressions above and below the spine that house important rotator cuff muscles.

THE UPPER LIMB: FROM SHOULDER TO FINGERTIPS

Each upper limb contains 30 bones. This section of the review sheet will focus on the humerus, radius, ulna, carpals, metacarpals, and phalanges.

The Humerus: The Arm Bone

The humerus is the long bone of the arm, extending from the shoulder to the elbow. Important features include the head, which fits into the glenoid cavity;

the greater and lesser tubercles, which are muscle attachment sites; and the deltoid tuberosity, a rough area on the shaft for the deltoid muscle. At the distal end, you will find the trochlea (which articulates with the ulna), the capitulum (which articulates with the radius), and the medial and lateral epicondyles, which are common sites for tendinitis, often referred to as "tennis elbow" or "golfer's elbow."

The Radius and Ulna: The

Forearm

The forearm consists of two parallel bones. In the anatomical position (palms facing forward), the radius is on the lateral side (thumb side), and the ulna is on the medial side (pinky side). This is a critical distinction for **Review Sheet Exercise 9 The Appendicular Skeleton Answers**.

- **The Ulna:** This bone is longer than the radius. Its prominent proximal feature is the olecranon process, which forms the point of the elbow. The trochlear notch lies inferior to the olecranon and articulates with the humerus. Distally, the ulna ends in a styloid process.

- **The Radius:** The radius is the primary bone involved in the wrist joint. Its proximal head is disc-shaped and rotates against the capitulum of the humerus. The radial tuberosity is a key muscle attachment point. Distally, the radius widens and features its own styloid process. The interosseous membrane connects the radius and ulna along their shafts.

The Wrist and Hand: Carpals,

Metacarpals, and Phalanges

The hand is a complex structure of 27 bones. The carpus (wrist) consists of eight small carpal bones arranged in two rows of four. A common mnemonic to remember them is "Some Lovers Try Positions That They Can't Handle" (Scaphoid, Lunate, Triquetrum, Pisiform, Trapezium, Trapezoid, Capitate, Hamate).

The palm of the hand is formed by five metacarpal bones, numbered 1 to 5 starting from the thumb side. Finally, the fingers contain 14 phalanges. Each finger (except the thumb) has three phalanges: proximal, middle, and distal.

The thumb has only two: proximal and distal. Knowing this arrangement is essential for any comprehensive review.

THE PELVIC GIRDLE: A STABLE FOUNDATION

Unlike the mobile pectoral girdle, the pelvic girdle is designed for stability and weight-bearing. It is formed by two coxal bones (hip bones) that articulate anteriorly at the pubic symphysis and posteriorly with the sacrum. Each coxal bone is actually a fusion of three separate bones: the ilium, ischium, and pubis. These bones fuse at the acetabulum, the deep socket that receives the head of the femur.

The Ilium, Ischium, and Pubis

These three bones are distinct in the developing skeleton but fuse by adulthood.

- **The Ilium:** The largest and most superior part of the coxal bone. Its superior margin is the iliac crest. The most superior point is the anterior superior iliac spine (ASIS), a common anatomical landmark.
- **The Ischium:** The strongest part of the hip bone. It forms the posterior-inferior portion. When

you sit, you are sitting on your ischial tuberosities. The ischial spine is a sharp projection that separates the greater and lesser sciatic notches.

- **The Pubis:** The anterior-inferior part. The two pubic bones meet at the pubic symphysis (a cartilaginous joint). The pubic arch is formed below the symphysis.

A key topic in **Review Sheet Exercise 9 The Appendicular Skeleton Answers** is often the comparison between a male and female pelvis. The female pelvis is generally broader, lighter, and has a wider pelvic inlet and outlet to facilitate childbirth. The pubic arch in females is wider (greater than 80 degrees), while in males it is more acute

(less than 70 degrees).

THE LOWER LIMB: DESIGNED FOR SUPPORT AND LOCOMOTION

The lower limbs are built for strength and stability. Each lower limb contains 30 bones, mirroring the upper limb in basic layout but differing greatly in form and function.

The Femur: The Thigh Bone

The femur is the longest, strongest, and heaviest bone in the body. Its proximal end features a spherical head that articulates with the acetabulum. The neck of the femur is a common site of

fracture in the elderly. Below the neck are the greater and lesser trochanters, large projections for powerful muscle attachments. Running down the posterior shaft is the linea aspera, a rough ridge for muscle attachment. Distally, the femur ends in the medial and lateral condyles, which articulate with the tibia.

The Patella: The Kneecap

The patella is a triangular sesamoid bone embedded within the quadriceps tendon. It protects the knee joint anteriorly and increases the leverage of the quadriceps muscle. It articulates with the patellar surface of the femur.

The Tibia and Fibula: The Leg

Like the forearm, the leg has two bones. The tibia is the weight-bearing bone (medial), and the fibula is a slender, non-weight-bearing bone (lateral).

- **The Tibia:** The shinbone. The proximal tibia features the medial and lateral condyles, which articulate with the femur. The tibial tuberosity is the bump below the kneecap where the patellar ligament attaches. The anterior border (shin) is a sharp ridge. The distal end forms the medial malleolus (the inner ankle bump).

- **The Fibula:** This bone does not articulate with the femur. Its proximal head is a small knob that articulates with the tibia. The distal end forms the lateral malleolus (the outer ankle bump). The fibula serves primarily as an attachment site for muscles.

The Ankle and Foot: Tarsals, Metatarsals, and

Phalanges

The foot contains 26 bones. It is designed to support body weight and act as a lever for walking. The tarsus (ankle) consists of seven tarsal bones. The largest is the calcaneus (heel bone). The talus sits on top of the calcaneus and articulates with the tibia and fibula to form the ankle joint. The other tarsals include the navicular, cuboid, and three cuneiform bones.

The five metatarsal bones form the instep of the foot. The phalanges of the toes follow the same pattern as the fingers: the great toe (hallux) has two phalanges, while the other four toes have three each. The foot also features several arches (medial longitudinal,

lateral longitudinal, and transverse) that distribute weight and absorb shock.

COMMON QUESTIONS AND CONCEPTS IN REVIEW SHEET EXERCISE 9

When you are working through **Review Sheet Exercise 9 The Appendicular Skeleton Answers**, you will likely encounter questions that go beyond simple identification. Here are some conceptual areas that are frequently tested:

Differences

Between the Pectoral and Pelvic Girdles

- The pectoral girdle does not directly articulate with the vertebral column (except via muscles and the clavicle), while the pelvic girdle is firmly attached to the sacrum.
- The pectoral girdle is optimized for mobility; the pelvic girdle is optimized for stability and weight-bearing.
- The socket for the upper limb (glenoid cavity) is shallow; the

socket for the lower limb (acetabulum) is deep.

Bone Surface Markings

You will need to distinguish between different types of bone markings. For example:

- **Depressions and openings:** Foramen (hole), fossa (shallow depression), sulcus (groove).
- **Processes that form joints:** Condyle (smooth, rounded projection), head (expanded end), facet (flat, smooth surface).
- **Processes that are attachment sites:** Tubercle (small, rounded

projection), tuberosity (large, rough projection), trochanter

(large, blunt projection, only on the femur), spine (sharp, slender