
Simple Probability Problems And Solutions

*Simple
Probability
Problems
And
Solutions*

*Downloaded from
www.everybodystreet.com
by Aidan & Turner*

UNDERSTANDING SIMPLE PROBABILITY PROBLEMS AND SOLUTIONS: A BEGINNER'S GUIDE

Probability is one of those fascinating branches of mathematics that touches nearly every part of our daily lives. Whether you are deciding whether to take an umbrella based on a weather forecast, playing a card game, or analyzing data for a

business report, probability is quietly at work. For many learners, the best way to build confidence is to start with **simple probability problems and solutions** that demonstrate core ideas without overwhelming complexity. This article walks through the fundamentals, offers clear examples, and shows you how to approach these problems step by step. By the end of this guide, you will have a solid grasp of basic probability concepts.

You will know how to define events, compute likelihoods, and avoid common pitfalls. And you will see exactly why mastering simple probability problems and solutions is the perfect foundation for more advanced statistical thinking.

WHAT IS PROBABILITY? A QUICK REFRESHER

At its heart, probability is a measure of how likely something is to happen. We express it as a number between 0 and 1, where 0 means the event is impossible and 1 means it is certain. You can also express probability as a percentage, a fraction, or a decimal.

The most basic formula for probability is:

Probability of an event = (Number of

**favorable outcomes)
/ (Total number of
possible outcomes)**

This formula assumes all outcomes are equally likely. This is a key point when working through simple probability problems and solutions. For example, when you flip a fair coin, the chance of getting heads is 1 out of 2, or $\frac{1}{2}$, or 0.5, or 50 percent.

KEY TERMS YOU NEED TO KNOW

Before diving into specific problems, let us clarify a few essential terms. These will appear again and again in simple probability problems and solutions.

- **Experiment:**
Any action or process that produces a well-defined outcome.

Flipping a coin is an experiment.

- **Sample space:** The set of all possible outcomes of an experiment. For a coin flip, the sample space is {Heads, Tails}.
- **Event:** A specific outcome or group of outcomes you are interested in. For example, rolling an even number on a die.
- **Favorable outcomes:** The outcomes that match your event.
- **Random:** When every outcome has an equal chance of occurring (in a fair setting).

Understanding these terms makes it much easier to follow along

with examples and to create your own simple probability problems and solutions.

CLASSIC COIN TOSS PROBLEMS

Coin toss problems are the most common starting point for simple probability problems and solutions. They are easy to visualize and involve only two possible outcomes.

Problem 1: What is the probability of

flipping the
heads probability
a fair coin of
getting
coin?

Solution: There is 1 favorable outcome (heads) out of 2 possible outcomes. So the probability is $1/2$ or 0.5 or 50 percent.

exactly
one
head?

Problem
2: If you
flip a coin
twice,
what is

Solution: First, list all possible outcomes for two flips: HH, HT, TH, TT. That is 4 outcomes. The favorable outcomes (exactly one head) are HT and TH. That is 2 favorable outcomes. So the probability is $2/4$, which simplifies to $1/2$ or 0.5.

Problem

3: What is the probability of getting at least one head in three coin flips?

Solution: The sample space for three flips has 8 outcomes ($2 \times 2 \times 2 = 8$). The only outcome without any heads is TTT. So the number of favorable outcomes is $8 - 1 = 7$.

The probability is $7/8$ or 0.875.

These simple probability problems and solutions show how listing outcomes and counting carefully gives you the answer.

DICE ROLLING PROBLEMS

A standard six-sided die is another classic tool for learning probability. Each face is equally likely to land facing up, assuming a fair die.

Problem

4: What is the probability

y of even
rolling a number?

4?

Solution: There is 1 favorable outcome (rolling a 4) out of 6 possible outcomes. Probability = $1/6 \approx 0.1667$.

Solution: The even numbers on a die are 2, 4, and 6. That is 3 favorable outcomes out of 6. Probability = $3/6 = 1/2 = 0.5$.

Problem

Problem

6: What

5: What

is the

is the

probabilit

probabilit

y of

y of

rolling a

rolling an

number

greater
than 4?

Solution: Numbers greater than 4 on a standard die are 5 and 6. That is 2 favorable outcomes out of 6. Probability = $2/6 = 1/3 \approx 0.3333$.

Problem
7: Roll
two dice.
What is
the
probabilit
y that the

sum is 7?

Solution: When rolling two dice, there are $6 \times 6 = 36$ total outcomes. The pairs that sum to 7 are: (1,6), (2,5), (3,4), (4,3), (5,2), (6,1). That is 6 favorable outcomes. Probability = $6/36 = 1/6 \approx 0.1667$.

These simple probability problems and solutions around dice are excellent for practicing counting and understanding sample spaces.

PROBLEMS INVOLVING MARBLES OR BALLS

Drawing marbles from a bag is a very common type of problem. These problems often introduce the idea of without replacement, which changes

probabilities.

Problem 8: A bag contains 3 red marbles, 2 blue marbles, and 5 green marbles. What is the

probability of drawing a red marble?

Solution: Total marbles = $3 + 2 + 5 = 10$. Favorable outcomes (red) = 3. Probability = $3/10 = 0.3$ or 30 percent.

Problem 9: From the same bag, what

is the
probability
of
drawing a
blue or
green

marble?

Solution: Blue
marbles = 2, green
marbles = 5. Favorable
outcomes = $2 + 5 = 7$.
Total = 10. Probability
= $7/10 = 0.7$.

Problem
10:

Without
replacem
ent, draw
two
marbles
from the
same
bag.

What is
the
probability
that

both are
red?

Solution: For the first draw, probability of red = $3/10$. After removing one red marble, there are 2 red marbles left and 9 total. So for the second draw, probability of red = $2/9$. Multiply the probabilities: $(3/10) \times (2/9) = 6/90 = 1/15 \approx 0.0667$.

The without replacement scenario is a critical variation in simple probability problems and solutions because it shows how events can depend on each other.

CARD PROBLEMS

A standard deck of 52 playing cards offers many opportunities for simple probability problems and

solutions. A deck has 4 suits (hearts, diamonds, clubs, spades), 13 cards per suit, and 26 red cards (hearts and diamonds) and 26 black cards (clubs and spades).

Problem 11: What is the probability of drawing a heart from a full deck?

Solution: There are 13 hearts in a deck of 52

cards. Probability =
 $13/52 = 1/4 = 0.25$.

Problem

Problem

13: What

12: What

is the

is the

probabilit

probabilit

y of

y of

drawing a

drawing a

red card

face card

that is

(Jack,

also a

Queen, or

face

King)?

card?

Solution: There are 4 suits, each with 3 face cards, so 12 face cards total. Probability = $12/52 = 3/13 \approx 0.2308$.

Solution: Red suits are hearts and diamonds. Each has 3 face cards, so 6 red face cards. Probability = $6/52 = 3/26 \approx$

0.1154.

Problem 14: Draw two cards without replacement. What is the probability both are aces?

Solution: There are 4 aces in the deck. First draw probability = $4/52$. After drawing one ace, 3 aces remain out of 51 cards. Second draw probability =

$3/51$. Multiply: $(4/52) \times (3/51) = 12/2652 = 1/221 \approx 0.0045$.

These card-based simple probability problems and solutions are great for practicing compound events and conditional probability.

PROBABILITY OF INDEPENDENT AND DEPENDENT EVENTS

Two events are independent if the outcome of one does not affect the outcome of the other. For example, coin flips are independent. Events are dependent if the outcome of one does affect the other, as we saw with drawing without replacement.

For independent events, multiply the individual probabilities: $P(A \text{ and } B) = P(A) \times P(B)$.

For dependent events, you adjust the second probability based on the first outcome.

Problem 15: Flip a coin and roll a die.

What is the probability of getting heads

and rolling a 3?

Solution: The events are independent.
 $P(\text{heads}) = 1/2$.
 $P(\text{rolling a 3}) = 1/6$.
Multiply: $(1/2) \times (1/6) = 1/12 \approx 0.0833$.

Problem 16: From a bag with 4 white and 6 black

marbles,
draw two
without
replacem
ent. What
is the
probab
y the f
is white
and the
second is
black?

marble, 9 marbles remain: 3 white and 6 black. Second draw: $P(\text{black}) = 6/9 = 2/3$. Multiply: $(2/5) \times (2/3) = 4/15 \approx 0.2667$.

These simple probability problems and solutions help distinguish between independent and dependent events, a crucial skill for moving forward.

USING THE COMPLEMENT RULE

Sometimes it is easier to find the probability of an event happening by finding the probability of it not happening and subtracting from 1. This is called the complement rule: $P(A) = 1 - P(\text{not } A)$.

Solution: First draw:
 $P(\text{white}) = 4/10 = 2/5$.
After drawing a white

Problem

17: What is the probability of rolling at least one 6 in two dice rolls?

Solution: It is easier to find the probability of rolling no 6 at all. For one die, $P(\text{not } 6) = 5/6$. For two independent rolls, $P(\text{no } 6) = (5/6) \times (5/6) = 25/36$. Then $P(\text{at least one } 6) = 1 - 25/36 = 11/36 \approx 0.3056$.

The complement rule is a powerful shortcut in many simple probability problems and solutions.

PROBABILITY IN EVERYDAY LIFE

Simple probability problems and solutions are not just for math class. They appear all around you. Weather forecasts use probability to express the chance of rain. Insurance companies use probability to set premiums. Game designers use probability to balance gameplay. Even doctors use probability to interpret test results.

Understanding basic probability helps you make better decisions. When you know the odds, you can evaluate risks more clearly. For example, if a medical

test for a rare disease is 99 percent accurate, the probability that you actually have the disease after a positive result is much lower than you might think. That is because the base rate (how rare the disease is) matters. This is a more advanced application