
6 4 Elimination Using Multiplication Practice And

6 4
Elimination
Using
Multiplication Practice And
Downloaded from
www.everybodystreet.com
by Brady & Phelps

INTRODUCTION: MASTERING THE ELIMINATION METHOD WITH MULTIPLICATION

Algebra often feels like a puzzle where each new technique unlocks a broader set of solutions. Among the most powerful tools in solving systems of linear equations is the **elimination method**, and when you add multiplication to the mix, you gain the ability to solve almost

any pair of standard linear equations. The section commonly labeled "**6 4 Elimination Using Multiplication Practice And**" in many algebra textbooks is specifically designed to help students move beyond simple elimination where coefficients already match. Instead, it teaches you how to strategically multiply one or both equations so that a variable cancels perfectly. This article will walk you through the concept,

step-by-step procedures, and ample practice to ensure you not only understand the method but can apply it confidently in homework, tests, and real-world problem solving.

Whether you are a student preparing for an exam, a parent helping with homework, or someone refreshing their math skills, mastering the elimination method with multiplication is a crucial milestone. By the end of this guide, you will have a clear roadmap for handling any system where the coefficients are not initially opposites or matches. We will break down the process, offer practice problems with solutions, and highlight common pitfalls to avoid. Let's dive into the world of **6 4**

Elimination Using Multiplication Practice And - your gateway to algebraic fluency.

UNDERSTANDING THE ELIMINATION METHOD

The elimination method is a technique for solving systems of two linear equations in two variables. The core idea is to combine the two equations in a way that eliminates one variable, allowing you to solve for the other. When the coefficients of one variable are already opposites or equal, you can simply add or subtract the equations. However, many systems do not come pre-arranged. That is where multiplication enters the picture.

For example, consider

the system:

- $3x + 2y = 12$
- $5x - 3y = 1$

Here, neither the x coefficients (3 and 5) nor the y coefficients (2 and -3) are opposites. If we try to add or subtract as is, we fail to eliminate a variable. By multiplying one or both equations by a carefully chosen number, we can create matching coefficients that cancel out. This technique is the heart of what is often taught under **6 4 Elimination Using Multiplication Practice And**.

When Multiplication Is

Necessary

Multiplication becomes necessary when the coefficients of the variable you wish to eliminate are not already negatives or equals. The general rule: you can multiply any equation by a nonzero constant without changing its solution set. Therefore, you can scale the equations to produce opposite coefficients for either x or y . The goal is to choose multipliers so that after addition or subtraction, one variable disappears.

Sometimes you need to multiply only one equation; other times both require scaling. For instance, to eliminate y in the

system above, you could multiply the first equation by 3 and the second by 2, giving you $6y$ and $-6y$. That would allow cancellation by addition. The **6 4 Elimination Using Multiplication Practice And** lessons typically cover both scenarios: multiplying one equation and multiplying both.

STEP-BY-STEP GUIDE FOR 6-4 ELIMINATION USING MULTIPLICATION

Following a systematic process ensures accuracy and builds intuition. Below are the steps you should use when solving a system with the elimination method that requires multiplication.

Step 1: Align the Equations

Write both equations in standard form: $Ax + By = C$. If an equation is given in slope-intercept form ($y = mx + b$), convert it to standard form first. Align the variables vertically so you can easily compare coefficients.

Example:

- $4x + 3y = 10$
- $2x - 5y = -8$

Step 2: Decide Which

to Variable Eliminate Step 3: Determine the Multiplier (s)

Choose the variable that will be easiest to eliminate. Look at the coefficients. Often it is easier to eliminate the variable whose coefficients have a smaller least common multiple (LCM) or that require smaller multipliers. In the example above, the x coefficients are 4 and 2; the y coefficients are 3 and -5. The LCM for x is 4 (since 4 and 2 align nicely), while for y the LCM is 15. So eliminating x may be simpler.

You need to multiply one or both equations so that the coefficients of the chosen variable become opposites. For our example, if we want to eliminate x , we can multiply the second equation by -2 (or the first by 1 and the second by -2). Multiplying the second equation ($2x - 5y = -8$) by -2 gives $-4x + 10y = 16$. Now the x coefficients are 4 and -4.

If you prefer to eliminate y , you would need to find multipliers

that make the y coefficients opposites. Multiply the first equation by 5 (giving $20x + 15y = 50$) and the second equation by 3 (giving $6x - 15y = -24$). Now y has coefficients $+15$ and -15 .

Tip: Choose the multipliers that result in the smallest possible numbers to simplify later arithmetic. This is a key point in **6 4 Elimination Using Multiplication Practice And** exercises.

Step 4:
Multiply
the

Equations

Perform the multiplication carefully. Multiply every term on both sides of the equation by the chosen constant. Do not forget to multiply the constant term (the number on the right side of the equals sign).

Using our example to eliminate x :

- First equation remains: $4x + 3y = 10$
- Second equation multiplied by -2 : $-4x + 10y = 16$

Step 5:
Add (or

the Subtract) Back Equations e to Find

Add the two equations together. Because the coefficients of x are now opposites ($4x$ and $-4x$), they cancel. If you had created equal coefficients instead of opposites, you would subtract. In this case, addition works.

$$\begin{array}{r} 4x + 3y = 10 \\ -4x + 10y = 16 \\ \hline \end{array}$$

$$0x + 13y = 26$$

So $13y = 26$, which gives $y = 2$.

Step 6:

the Second Variable

Take the value you found ($y = 2$) and substitute it into one of the original equations (preferably one with simple numbers) to solve for x .

$$\begin{aligned} \text{Using } 4x + 3(2) &= 10 \rightarrow \\ 4x + 6 &= 10 \rightarrow 4x = 4 \\ \rightarrow x &= 1. \end{aligned}$$

Thus the solution is $(1, 2)$. Always check by plugging into the other original equation: $2(1) - 5(2) = 2 - 10 = -8$, which matches.

PRACTICE PROBLEMS FOR 6-4 ELIMINATION USING MULTIPLICATION

To truly internalize the method, work through a variety of problems. Below are three practice problems that reflect typical **6 4 Elimination Using Multiplication Practice And** exercises. Try them on your own before reading the solutions.

Problem 1

Solve the system:

- $3x + 4y = 19$
- $5x - 2y = 17$

Solution: Choose to eliminate y . Multiply the first equation by 1 and the second

equation by 2 to get y coefficients of 4 and -4 .

- $3x + 4y = 19$
(unchanged)
- Multiply second equation by 2:
 $10x - 4y = 34$

Add: $13x = 53 \rightarrow x = 53/13 \approx 4.0769$ (or keep as fraction $53/13$ if exact). Actually, $53/13$ simplifies? 53 is prime, 13 is prime, so $53/13$. Then substitute to find y : $3(53/13) + 4y = 19 \rightarrow 159/13 + 4y = 247/13 \rightarrow 4y = (247-159)/13 = 88/13 \rightarrow y = 22/13$. So solution $(53/13, 22/13)$.

Problem 2

Solve:

- $2x + 3y = 8$
- $4x - 5y = -6$

Solution: Eliminate x by multiplying the first equation by -2 (so that $2x$ becomes $-4x$).

- Multiply first by -2 : $-4x - 6y = -16$
- Second equation: $4x - 5y = -6$

Add: $-11y = -22 \rightarrow y = 2$. Substitute into first original: $2x + 3(2) = 8 \rightarrow 2x + 6 = 8 \rightarrow 2x = 2 \rightarrow x = 1$. Solution $(1, 2)$.

Problem 3

Solve (requires multiplying both equations):

- $7x + 2y = 13$
- $3x - 5y = 8$

Solution: To eliminate y , multiply the first equation by 5 and the second by 2 to get $10y$ and $-10y$.

- $5(7x + 2y = 13) \rightarrow 35x + 10y = 65$
- $2(3x - 5y = 8) \rightarrow 6x - 10y = 16$

Add: $41x = 81 \rightarrow x = 81/41$. Substitute into first original: $7(81/41) + 2y = 13 \rightarrow 567/41 + 2y = 533/41 \rightarrow 2y = (533 - 567)/41 = -34/41 \rightarrow y = -17/41$. Solution $(81/41, -17/41)$.

COMMON MISTAKES TO AVOID IN ELIMINATION USING MULTIPLICATION

Even experienced students make errors. Here are several pitfalls that the **6 4 Elimination Using Multiplication Practice And** materials often emphasize:

- **Forgetting to multiply every**

term: When you multiply an equation, you

must multiply the constant term on the right side as well