

Introduction To Lagrangian And Hamiltonian Mechanics

Introduction To Lagrangian And Hamiltonian Mechanics Downloaded from www.everybodystreet.com by Riley & Johns

INTRODUCTION TO LAGRANGIAN AND HAMILTONIAN MECHANICS: A FRESH PERSPECTIVE ON MOTION

When students first encounter classical mechanics, they typically study Newton's laws of motion. This approach, while powerful, often becomes cumbersome when dealing with complex systems like pendulums with moving supports, rotating bodies, or multi-particle interactions. Enter Lagrangian and Hamiltonian mechanics — two reformulations of classical physics that provide deeper insight and greater mathematical elegance. This introduction to Lagrangian and Hamiltonian mechanics will explore how these frameworks simplify problem-solving, reveal conserved quantities, and lay the groundwork for modern physics, including quantum mechanics and relativity.

Both Lagrangian and Hamiltonian mechanics shift focus from forces to energies. Where Newton uses vectors and forces, these methods use scalar quantities — kinetic and potential energy — and exploit symmetry to find conservation laws. This makes them indispensable for advanced physics and engineering. Let us begin our journey by understanding the foundational concepts and then seeing how they interconnect.

THE CORE IDEA: ENERGY OVER FORCE

Newton's second law, $\mathbf{F} = m\mathbf{a}$, requires knowing every force acting on a system. In many problems, forces are difficult to model directly (e.g., constraints like a bead on a wire). Lagrangian mechanics bypasses this by introducing the **Lagrangian** function, usually denoted as $L = T - V$, where T is kinetic energy and V is potential energy. The equations of motion are then derived from a single principle: the principle of least action.

Similarly, Hamiltonian mechanics introduces the **Hamiltonian** ($H = T + V$) (total mechanical energy in many cases) and uses it to generate equations of motion that are first-order in time, offering a symmetrical structure that is ideal for connecting with quantum theory.

LAGRANGIAN MECHANICS: THE PRINCIPLE OF LEAST ACTION

The Action Functional

The centerpiece of Lagrangian mechanics is the **action** S , defined as the time integral of the Lagrangian along a path:

$$S[q(t)] = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt$$

Here, q represents generalized coordinates — any set of variables that uniquely describe the system's configuration (e.g., angles, distances, or even more abstract quantities). The Principle of Least Action (or Hamilton's principle) states that the actual path taken by the system between two times minimizes (or extremizes) the action. This leads to the Euler-Lagrange equations:

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_i} \right) - \frac{\partial L}{\partial q_i} = 0$$

For each generalized coordinate q_i , these equations yield the equations of motion without needing to account for constraint forces explicitly.

Advantages of the Lagrangian Formulation

- Coordinate independence:** You can choose any generalized coordinates that fit the problem — even non-inertial ones — and the Euler-Lagrange equations remain valid.
- Handling constraints:** Holonomic constraints (relationships between coordinates) can be incorporated via Lagrange multipliers or by reducing the number of coordinates.
- Symmetry and conservation:** If the Lagrangian does not depend explicitly on a coordinate q_i , then the conjugate momentum $p_i = \frac{\partial L}{\partial \dot{q}_i}$ is conserved (Noether's theorem).

A Simple Example: The Simple Pendulum

For a pendulum of length l and mass m swinging in a plane, the natural coordinate is the angle θ . The kinetic energy is $T = \frac{1}{2} m l^2 \dot{\theta}^2$ and potential energy is $V = mgl(1 - \cos\theta)$. The Lagrangian is:

$$L = T - V = \frac{1}{2} m l^2 \dot{\theta}^2 - mgl(1 - \cos\theta)$$

Applying the Euler-Lagrange equation gives:

$$m l^2 \ddot{\theta} + mgl \sin\theta = 0 \quad \text{or} \quad \ddot{\theta} + \frac{g}{l} \sin\theta = 0$$

This is the familiar equation for a pendulum, derived without worrying about tension forces. The

Lagrangian method elegantly yields the correct motion.

Degrees of Freedom and Generalized Coordinates

One of the most powerful aspects of Lagrangian mechanics is the reduction of degrees of freedom. A system of N particles in 3D space normally has $3N$ coordinates, but constraints reduce this number. For example, a rigid body has only 6 degrees of freedom (3 translation, 3 rotation). Generalized coordinates allow you to work with exactly the minimum number of variables needed.

HAMILTONIAN MECHANICS: FROM SECOND ORDER TO FIRST ORDER

The Legendre Transform

Hamiltonian mechanics arises from a mathematical transformation called the **Legendre transform**. Starting from the Lagrangian $L(q, \dot{q}, t)$, we define the conjugate momentum:

$$p_i = \frac{\partial L}{\partial \dot{q}_i}$$

Then the Hamiltonian H is defined as:

$$H(q, p, t) = \sum_i p_i \dot{q}_i - L$$

If the Lagrangian is not explicitly time-dependent and the constraints are scleronomic (time-independent), then H equals the total energy $T+V$. The Hamiltonian is a function of generalized coordinates and momenta, not velocities.

Hamilton's Equations

Instead of second-order differential equations (like the Euler-Lagrange equations), Hamilton's equations are a set of first-order equations:

$$\dot{q}_i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q_i}$$

These $2n$ equations (for n degrees of freedom) are symmetrical and often easier to solve numerically. They also have a beautiful geometric interpretation: the system's state moves along a trajectory in **phase space**, a space with coordinates (q_i, p_i) .

Phase Space and Flow

Phase space is a cornerstone of Hamiltonian mechanics. Each point corresponds to a unique state of the system. As time evolves, the point traces a curve. An important property is **Liouville's theorem**: the volume of a region of phase space remains constant under Hamiltonian flow. This is a fundamental result in statistical mechanics and ergodic theory.

Example: Harmonic Oscillator in Hamiltonian Form

For a one-dimensional harmonic oscillator with mass m and spring constant k , the Hamiltonian is:

$$H = \frac{p^2}{2m} + \frac{1}{2} k q^2$$

Hamilton's equations become:

$$\dot{q} = \frac{p}{m}, \quad \dot{p} = -k q$$

Differentiating the first and substituting the second gives $\ddot{q} + (k/m) q = 0$, which is the familiar simple harmonic motion equation. The first-order form is ideal for numerical integration.

COMPARING LAGRANGIAN AND HAMILTONIAN METHODS

Similarities

- Both rely on energy (scalars) rather than forces (vectors).
- Both are coordinate independent and can handle constraints elegantly.
- Both lead to conservation laws through symmetries.

Differences