

---

# How To Take Magnesium Citrate Oral Solution

---

*How To Take  
Magnesium Citrate Oral  
Solution*

*Downloaded from  
[www.everybodystreet.com](http://www.everybodystreet.com)  
by Guerra & Micah*

---

## UNDERSTANDING HOW TO TAKE MAGNESIUM CITRATE ORAL SOLUTION SAFELY AND EFFECTIVELY

---

Magnesium citrate oral solution is a widely used and effective remedy for relieving occasional constipation, preparing the bowel for medical procedures, and supporting overall

digestive health. While it is available over the counter and generally considered safe, knowing exactly **how to take magnesium citrate oral solution** correctly can make a significant difference in its effectiveness and your comfort. Many people underestimate the importance of proper dosing, timing, and hydration when using this laxative. This article provides a clear, step-by-step guide on using magnesium citrate oral solution, along

with essential safety information and practical tips to help you get the best results.

---

## WHAT IS MAGNESIUM CITRATE ORAL SOLUTION?

---

Magnesium citrate is a combination of magnesium and citric acid that works as an osmotic laxative. When you take the solution orally, it draws water into the intestines, which softens the stool and stimulates bowel movements. Unlike stimulant laxatives, magnesium citrate helps the body move waste along naturally without forcefully contracting the colon. This makes it a preferred option for many people seeking relief from constipation or needing to clear the bowel before a colonoscopy or other medical examination.

The solution is typically sold as a clear, fizzy liquid in a bottle. It has a distinct sour or lemony taste, which some people find unpleasant. Understanding the proper way to take it, including how to handle the taste and timing, is essential for a positive experience.

---

## BEFORE YOU BEGIN: IMPORTANT PRECAUTIONS

---

Before you learn **how to take magnesium citrate oral solution**, it is crucial to consider whether this product is suitable for you. Consult your doctor if you have any of the following conditions:

- Kidney disease or impaired kidney function
- Heart problems, including arrhythmias or a history of heart

**CITRATE ORAL SOLUTION: A**

- Stomach ulcers or other gastrointestinal disorders
- Electrolyte imbalances
- If you are pregnant, planning to become pregnant, or breastfeeding
- If you are on a magnesium-restricted diet

Additionally, magnesium citrate can interact with certain medications, including antibiotics, diuretics, and medications for osteoporosis or thyroid conditions. Always tell your healthcare provider about all medications and supplements you are currently taking.

---

**HOW TO TAKE MAGNESIUM**

---

**STEP-BY-STEP GUIDE**

---

## Step 1: Read the Label Carefully

Before opening the bottle, read the entire label and the package insert if one is included. Different brands may have slightly different concentrations or instructions. The label will tell you the exact dosage, how often you can take it, and any specific warnings. Pay close attention if you are using the solution for colonoscopy preparation, as the instructions may differ from those for general constipation relief.

## Step 2: Determine the Correct Dosage

Dosage recommendations vary based on age and the purpose of use. For adults with occasional constipation, the typical dose is one full bottle (10 fluid ounces or 300 mL) taken once daily, or as directed by a doctor. Do not exceed the recommended dose in a 24-hour period. For children, the dosage is usually smaller and should be determined by a pediatrician. Never guess the dosage for a child.

## Step 3: Chill the Solution for Better Taste

One of the most common complaints about magnesium citrate is its taste. The solution is sour and can be somewhat unpleasant. To make it more palatable, refrigerate the bottle for at least one hour before drinking. A cold solution is much easier to tolerate. You can also add a small amount of ice to the glass after pouring, but avoid diluting it with too much liquid, as this may reduce its effectiveness.

## Step 4: Take the Solution on an Empty Stomach

For best results, take magnesium citrate oral solution on an empty stomach. Food can interfere with its absorption and delay its laxative effect. Plan to take it at least two hours after your last meal. Many people find it convenient to take it first thing in the morning or a few hours after lunch.

## Step 5: Drink a Full Glass of Water Afterward

Immediately after taking the recommended dose, drink a full 8-ounce glass of water or clear liquid. This step is critical because magnesium citrate works by drawing water into the bowel. If you are dehydrated, the solution will not work as effectively, and you may experience cramping or discomfort. Staying well-hydrated throughout the day enhances the laxative effect and helps prevent electrolyte imbalances.

## Step 6: Stay Near a Bathroom

Magnesium citrate typically produces a bowel movement within 30 minutes to 6 hours after taking it. The timing can vary from person to person. It is wise to remain near a bathroom for several hours after taking the solution. Do not plan any activities that would make it difficult to access a restroom quickly.

## Step 7: Do Not

## Use for More Than 7 Consecutive Days

Magnesium citrate is intended for short-term use only. Using it for more than seven consecutive days can lead to dependence on laxatives, electrolyte imbalances, and other health issues. If your constipation does not improve after one week, stop using the solution and consult your doctor.

---

**PROPER DOSAGE GUIDELINES**  
**FOR DIFFERENT AGE GROUPS**

| Age Group                   | Recommended Dosage                                     | Maximum Frequency        |
|-----------------------------|--------------------------------------------------------|--------------------------|
| Adults (18 years and older) | 10 fl oz (300 mL) once daily                           | Once per 24 hours        |
| Children (6 to 17 years)    | Consult a doctor (usually half the adult dose or less) | As directed by physician |
| Children under 6 years      | Do not use without medical advice                      | Not applicable           |

Dosage may vary depending on the specific brand and concentration. Always follow the instructions on the product you purchase. If you are using

dosage instructions for colonoscopy preparation, your doctor will give you specific timing and dosage instructions that may differ from the general guidelines above.

**TIPS FOR BEST RESULTS WHEN TAKING MAGNESIUM CITRATE**

Use a Straw to Minimize Taste

Drinking the chilled solution through a straw can help bypass the taste buds on your tongue, making the experience less unpleasant. Place the straw toward the back of your mouth and swallow quickly.

## Follow with a Sip of Juice or Ginger Ale

After taking the solution and drinking your glass of water, you can take a small sip of apple juice, clear broth, or ginger ale to help clear the taste from your mouth. Avoid citrus juices, as they can interact with the solution and cause additional sourness.

## Time Your Dose

## Carefully

If you are using magnesium citrate for constipation relief, take it at a time when you will be at home for several hours. Avoid taking it right before bed unless you are prepared for a potential middle-of-the-night bathroom visit. For colonoscopy prep, follow your doctor's timing instructions precisely.

## Avoid Alcohol and Caffeine

Alcohol and caffeine can dehydrate the body, which counteracts the effects of magnesium citrate. Stick with clear fluids like water, herbal tea, or clear broth



while the solution is working.

---

### **WHAT TO EXPECT AFTER TAKING MAGNESIUM CITRATE**

---

Within 30 minutes to 6 hours after taking the solution, you will likely experience a strong urge to have a bowel movement. The stool will be watery or semi-liquid. This is normal and indicates that the solution is working. You may experience some abdominal cramping, bloating, or gassiness. These side effects are usually mild and temporary.

For many people, a single dose is sufficient to relieve constipation. If you do not have a bowel movement within six hours, do not take another dose. Wait 24 hours before repeating the process, and consult your doctor if constipation persists.

---

### **POTENTIAL SIDE EFFECTS AND WHEN TO SEEK MEDICAL HELP**

---

## Common Side Effects

- Abdominal cramping or discomfort
- Bloating and gas
- Nausea or vomiting (rare)
- Diarrhea (expected effect)
- Dizziness or lightheadedness (if dehydrated)

## Serious Side

# Effects (Seek Medical Attention Immediately)

- Severe or persistent abdominal pain
- Rectal bleeding or bloody stools
- Signs of dehydration (extreme thirst, dry mouth, little or no urination, weakness)
- Irregular heartbeat or chest pain
- Muscle weakness or cramping
- Allergic reaction (rash, itching, swelling, difficulty breathing)

If you experience any serious side effects, stop using the product

immediately and contact your healthcare provider or seek emergency medical care.

---

## **WHO SHOULD AVOID MAGNESIUM CITRATE**

---

Certain individuals should not take magnesium citrate without medical supervision. This includes people with:

- Severe kidney disease or renal failure
- Intestinal blockage or perforation
- Acute abdominal pain of unknown cause
- Electrolyte disorders such as hypermagnesemia
- Heart block or other serious heart conditions
- Nausea or vomiting

- Appendicitis or symptoms of appendicitis

If you are unsure whether magnesium citrate is safe for you, it is always better to ask your doctor before using it.

---

### **FREQUENTLY ASKED QUESTIONS ABOUT MAGNESIUM CITRATE ORAL SOLUTION**

---

Can I mix  
magnesium  
citrate with

juice?

While you can mix the solution with a small amount of clear juice to improve taste, it is best to avoid mixing it with citrus juices or other acidic beverages. The added acidity can make the taste even more sour. Plain water or clear broth is the safest option.

How long does  
magnesium  
citrate take to

## work?

Most people experience a bowel movement within 30 minutes to 6 hours after taking the solution. If you do not have a bowel movement within six hours, do not take another dose without consulting your doctor.

## Can I take magnesium citrate every

## day?

No. Magnesium citrate is intended for short-term use only. Using it daily for more than seven days can lead to laxative dependence, electrolyte imbalances, and other health problems. If you have chronic constipation, talk to your doctor about a long-term management plan.

## Is magnesium citrate safe for

## colonoscopy prep?

Yes, magnesium citrate is commonly used as part of colonoscopy bowel preparation. However, you should follow your doctor's specific instructions carefully. You may be required to take the solution in split doses and combine it with other laxatives or a clear liquid diet.

## Can I take magnesium

## citrate if I am on antibiotics?

Magnesium can interfere with the absorption of certain antibiotics, including tetracyclines and fluoroquinolones. If you are taking antibiotics, ask your doctor or pharmacist if you should avoid magnesium citrate or take it at a different time.

## What should I

# eat after taking magnesium citrate?

After the laxative effect has passed and you feel comfortable, start with light, easy-to-digest foods such as toast, crackers, rice, or bananas. Avoid heavy, greasy, or spicy foods for a day to allow your digestive system to recover.

---

## CONCLUSION

---

Knowing **how to take magnesium citrate oral solution** properly is the key to using it safely and effectively. Whether you are dealing with occasional

constipation or preparing for a medical procedure, following the correct steps—reading the label, chilling the solution, taking it on an empty stomach, staying hydrated, and staying near a bathroom—can make the experience much more manageable. Always adhere to the recommended dosage, do not use the solution for more than seven consecutive days, and seek medical advice if you have any underlying health conditions or concerns. Magnesium citrate can be a reliable and gentle solution when used responsibly, but like any medication, it deserves respect and caution. If you have questions about your specific situation, your healthcare provider is your best resource for personalized advice.