
How To Make Your Vagina Smell Nice

*How To
Make Your
Vagina Smell
Nice*

*Downloaded from
www.everybodystreet.com
by Anastasia & Orlando*

INTRODUCTION: UNDERSTANDING VAGINAL TIGHTNESS AND PELVIC HEALTH

Vaginal tightness is a topic that many women think about but rarely discuss openly. Whether due to childbirth, aging, hormonal changes, or simply a desire to enhance intimacy, the question "How to make your vagina tighter" is both common and valid. However, it is essential to approach this subject with accurate information,

respect for your body, and a focus on overall pelvic health. This article provides a comprehensive, evidence-based guide that covers natural exercises, lifestyle adjustments, medical options, and the most important thing of all—understanding what "tightness" really means for your body and your well-being.

Before exploring methods, let us clarify one key point: a vagina is not a muscle that needs to be "tightened" in the same way a fist clenches. Rather, vaginal tone is largely

determined by the strength of the pelvic floor muscles. These muscles support the bladder, uterus, and rectum, and they play a crucial role in sexual sensation, urinary control, and core stability. When these muscles weaken, a feeling of looseness may occur. Strengthening them is the most natural and effective way to achieve a tighter sensation for both you and your partner.

WHAT CAUSES A FEELING OF VAGINAL LOOSENESS?

Understanding the root causes helps you choose the right approach. Common factors that contribute to decreased vaginal tone include:

- **Childbirth:** Vaginal delivery stretches the pelvic floor muscles and the vaginal walls. While they recover significantly, some women experience persistent laxity.
- **Aging and Menopause:** Lower estrogen levels reduce tissue elasticity and muscle strength, leading to thinning and loosening of vaginal tissues.
- **Genetics:** Some women naturally have less muscle tone or more elastic connective tissue.
- **Weight Changes:** Significant

weight gain or loss can affect pelvic floor support.

- **Chronic**

Straining:

Constipation, heavy lifting, or chronic coughing can weaken the pelvic floor over time.

- **Lack of Use:**

Like any muscles, if pelvic floor muscles are not used regularly, they can become weaker.

It is also important to note that perceived looseness may be related to arousal or lack thereof. During sexual arousal, the vagina naturally expands and lubricates. If you are not fully aroused, the vagina may feel tighter, but the goal

should not be to mimic that state artificially. Instead, focus on overall pelvic health and strength.

NATURAL AND NON-INVASIVE METHODS

Pelvic Floor Exercises (Kegels)

Kegel exercises are the most well-known method for improving vaginal tightness. They target the pubococcygeus muscle, part of the pelvic floor. When done correctly and consistently, Kegels can significantly increase muscle tone and control.

How to perform Kegels correctly:

1. Identify the correct muscles:
The next time you urinate, try to stop the flow midstream. The muscles you use to do that are your pelvic floor muscles. Do not make a habit of stopping urine flow—only use this method to locate them.
2. Empty your bladder fully before exercising.
3. Sit or lie down in a comfortable position.
Squeeze your pelvic floor muscles as if you are trying to hold in urine or gas.
4. Hold the squeeze for 3-5 seconds, then relax

completely for 3-5 seconds.

5. Repeat 10 times, three times a day. Over time, increase the hold to 10 seconds.

Tips for best results:

- Breathe freely; do not hold your breath.
- Keep your abdomen, thighs, and buttocks relaxed.
- Consistency is key—you should see improvement in 4-6 weeks.

If you are unsure whether you are doing them correctly, a pelvic floor physical therapist can provide biofeedback or use an electronic device to confirm muscle activation.

Kegel Weighted Cones and Vaginal Balls

For women who want to add resistance to their Kegel routine, weighted devices such as cones or ben wa balls can be very effective. These are inserted into the vagina, and the natural urge to hold them in place activates the pelvic floor muscles. Start with a lighter weight and gradually increase as your muscles strengthen.

Using vaginal

weights safely:

- Always clean the device with warm soapy water before and after use.
- Use a water-based lubricant for comfort.
- Start with 10–15 minutes daily, while walking or doing household chores.
- If you feel any pain or discomfort, discontinue use and consult a healthcare provider.

Pelvic Floor Physical

Therapy

Pelvic floor physical therapy (PFPT) is a highly specialized and effective treatment. A trained therapist can assess your individual muscle strength, coordination, and any tightness or trigger points that may actually be causing pain rather than looseness. They use hands-on techniques, biofeedback, and personalized exercise programs to help you regain tone and function.

PFPT is particularly beneficial for women who:

- Have had a difficult childbirth or episiotomy.
- Experience urinary incontinence.

- Feel that their Kegel exercises are not working.
- Have pelvic organ prolapse.

Yoga and Pilates for Pelvic Strength

Yoga and Pilates can enhance pelvic floor awareness and strength indirectly through core stabilization postures. Specific poses like "Bridge Pose" (Setu Bandhasana), "Happy Baby" (Ananda Balasana), and "Cat-Cow" (Marjaryasana-Bitilasana) engage the pelvic floor in a controlled manner. Pilates focuses on the "Powerhouse," which

includes the pelvic floor, deep abdominals, and back muscles. Regular practice can improve posture, breath coordination, and overall muscular support.

LIFESTYLE AND DIETARY ADJUSTMENTS

Hydration and Fiber Intake

Chronic constipation forces you to strain during bowel movements, which can weaken the pelvic floor over time. Eating a high-fiber diet (fruits, vegetables, whole grains, legumes) and staying well-hydrated helps keep stools soft, reducing straining.

Adequate hydration also maintains vaginal tissue elasticity and lubrication.

Healthy Weight Managem ent

Excess body weight places additional pressure on the pelvic floor, contributing to muscle fatigue and laxity. Conversely, losing weight through a balanced diet and regular exercise can relieve some of that pressure and improve overall muscle tone. Avoid extreme or rapid weight loss, as that can also affect muscle mass and skin elasticity.

Hormonal Balance

Estrogen plays a vital role in maintaining vaginal tissue thickness, elasticity, and lubrication. During menopause or after certain medical treatments, estrogen levels drop, leading to vaginal atrophy. If you suspect hormonal imbalance is contributing to a feeling of looseness, consult a gynecologist. Options like vaginal estrogen creams, rings, or low-dose systemic therapy can restore tissue health. However, do not self-medicate; always seek professional guidance.

MEDICAL AND COSMETIC

OPTIONS

If natural methods prove insufficient or if the looseness significantly impacts your quality of life, several medical procedures exist. These should be considered only after a thorough evaluation by a qualified healthcare professional.

Laser Vaginal Rejuvenation

Laser treatments (such as FotonaSmooth or MonaLisa Touch) use thermal energy to stimulate collagen production and tighten the vaginal lining. The procedure is typically

performed in a clinic, requires no incisions, and has minimal downtime. It can also improve lubrication and reduce dryness. Results can last up to 12–18 months, and maintenance sessions may be needed.

Important considerations:

- These procedures are not FDA-approved specifically for vaginal "tightening," though they are used off-label.
- Not all women are candidates—a thorough pelvic exam is necessary.
- Cost can be high and is not usually covered by insurance.

Radiofrequency Treatments

Similar to laser, radiofrequency devices heat the tissue to promote collagen remodeling.

Treatments like ThermiVa or Viveve are non-invasive and done in a series of sessions. They can improve vaginal laxity, stress urinary incontinence, and overall comfort.

Vaginoplasty and

Perineoplasty

Surgical interventions involve tightening the vaginal muscles and the surrounding skin. Vaginoplasty narrows the vaginal canal, while perineoplasty tightens the perineum (the area between the vagina and anus). These are typically performed under anesthesia and require several weeks of recovery. They are most often chosen by women who have significant laxity after childbirth or aging, and who have not found relief through conservative measures.

Risks include infection, scarring, discomfort during intercourse, and potential changes in

sensation. Always choose a board-certified gynecologic surgeon with significant experience.

NATURAL SUPPLEMENTS AND HERBAL REMEDIES

Some women turn to herbal supplements marketed for vaginal tightness, such as "tightening" creams or pills containing astringents like alum, witch hazel, or plant extracts. The evidence for their effectiveness is minimal, and many can irritate sensitive tissues or disrupt the natural vaginal microbiome. **Using astringent products internally is not recommended** without medical supervision, as they can cause inflammation, dryness,

or even burns. A healthy, well-hydrated vaginal lining is far more beneficial than artificially constricted tissues.

THE DOS AND DON'TS OF VAGINAL TIGHTENING

Avoid common pitfalls that can harm your pelvic health:

- **Don't** overdo Kegel exercises. Just like any muscle, the pelvic floor can become too tight (hypertonic), leading to pelvic pain, painful intercourse, or difficulty urinating. Balance strength with relaxation.
- **Don't** use harsh chemical products or douches in an attempt to tighten the vagina. They disrupt pH and natural flora, increasing the risk of infections like bacterial vaginosis or yeast overgrowth.
- **Don't** ignore pain. If an exercise or treatment causes discomfort, stop and consult a professional.
- **Do** combine exercises with relaxation techniques. A healthy pelvic floor is both strong and flexible.
- **Do** maintain regular sexual activity or self-stimulation, as this naturally

increases blood flow and maintains vaginal health.

- **Do** communicate with your healthcare provider about any concerns. Many women suffer in silence, but solutions exist.

CONCLUSION: EMBRACE PELVIC HEALTH OVER PERFECTION

The question "How to make your vagina tighter" often arises from a desire to improve sexual satisfaction or to regain a sense of youthfulness. Yet true vaginal health is not about achieving a

specific tightness, but about having strong, responsive, and well-supported pelvic floor muscles. Start with non-invasive methods like Kegel exercises, pelvic floor therapy, and lifestyle changes. These are safe, effective, and empower you with body awareness. If you still feel that something is lacking, consult a specialist who can offer advanced options tailored to your unique anatomy and goals. Remember, every body changes over time—and that is perfectly normal. The most important step is to approach the topic with patience, knowledge, and self-compassion.