

# Solubility Curve Worksheet With Answers

Solubility Curve Worksheet With Answers

Downloaded from [www.everybodystreet.com](http://www.everybodystreet.com) by Siena & Kendall

## MASTERING CHEMISTRY: A COMPLETE GUIDE TO THE SOLUBILITY CURVE WORKSHEET WITH ANSWERS

For many students, the journey through high school or introductory college chemistry presents a series of challenging concepts. Among these, understanding how substances dissolve in a solvent under varying temperatures is a fundamental skill. The primary tool used to visualize and analyze this relationship is the solubility curve. Consequently, the **solubility curve worksheet with answers** has become a staple resource in classrooms and home study environments. This article aims to be your definitive guide, transforming a potentially confusing worksheet into a clear, manageable, and even enjoyable learning experience. We will explore what these graphs represent, how to interpret them, and how to tackle the common questions found on a **solubility curve worksheet with answers**, ensuring you build a solid foundation in solution chemistry.

Before we dive into the specifics of worksheets, it is essential to grasp why understanding solubility curves matters. Solubility is not a fixed property for most solids; it typically changes with temperature. Imagine trying to sweeten iced tea with sugar compared to hot tea. The hot tea can dissolve significantly more sugar. A solubility curve quantifies this relationship, allowing chemists to predict how much of a substance can be dissolved at a given temperature. This has real-world applications, from formulating medications and creating fertilizers to understanding geological processes like the formation of limestone caves. Therefore, mastering the **solubility curve worksheet with answers** is not just about passing a test; it is about understanding a principle that governs countless natural and industrial processes.

## WHAT EXACTLY IS A SOLUBILITY CURVE?

A solubility curve is a graphical representation of the solubility of a solute in a solvent as a function of temperature. Typically, the graph plots temperature on the x-axis (horizontal) and solubility on the y-axis (vertical). Solubility is most often expressed as grams of solute per 100 grams of water (g/100g H<sub>2</sub>O) or, less commonly, as grams per liter (g/L). Each line on the graph represents a different chemical substance, such as potassium nitrate (KNO<sub>3</sub>), sodium chloride (NaCl), or cerium sulfate (Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>).

The distinct shape of each line tells a story. Most solid solutes, like sugar and many salts, show an increase in solubility as the temperature rises. This is represented by an upward-sloping line. However, some substances are nearly flat, indicating their solubility changes very little with temperature (sodium chloride is a classic example). A few rare substances even have a downward slope, meaning their solubility decreases as the temperature goes up. Understanding these patterns is the first step in successfully using a **solubility curve worksheet with answers**.

## Key Terminology for Your Worksheet

To effectively work through any **solubility curve worksheet with answers**, you need to be comfortable with a few core terms. These words are frequently used in the questions and instructions.

- **Solute:** The substance being dissolved (the powder or solid on the curve).
- **Solvent:** The substance doing the dissolving (in our case, almost always water).
- **Solution:** A homogeneous mixture of solute and solvent.
- **Saturated Solution:** A solution that contains the maximum amount of solute that can be dissolved at a given temperature. On the graph, this is any point that lies exactly on the line of the curve.
- **Unsaturated Solution:** A solution that can still dissolve more solute. On the graph, this is any point that lies below the line of the curve.
- **Supersaturated Solution:** A highly unstable solution that contains more solute than is normally possible at a given temperature. On the graph, this is any point that lies above the line of the curve.
- **Solubility:** The maximum amount of solute that can be dissolved in a given amount of solvent (usually 100g of water) at a specific temperature to create a saturated solution.

## NAVIGATING YOUR SOLUBILITY CURVE WORKSHEET WITH ANSWERS: A STEP-BY-STEP GUIDE

A typical **solubility curve worksheet with answers** will present you with two main components: a graph and a series of questions. Let us break down the common question types and the logical steps to solve them.

## Question Type 1: Reading Solubility at a Specific Temperature

This is the most fundamental skill. The question will ask, "What is the solubility of potassium nitrate (KNO<sub>3</sub>) at 40°C?"

**How to solve it:**

1. Find the temperature (40°C) on the x-axis.

2. Draw an imaginary vertical line straight up from that point until it hits the curve for the specific substance (e.g., the KNO<sub>3</sub> curve).
3. From that intersection point, draw an imaginary horizontal line across to the y-axis.
4. Read the value on the y-axis. For KNO<sub>3</sub> at 40°C, this is typically around 64 g/100g H<sub>2</sub>O.

**Answer on the worksheet:** 64 g/100g H<sub>2</sub>O. This means that to make a saturated solution at 40°C, you would need to dissolve 64 grams of KNO<sub>3</sub> in 100 grams of water.

## Question Type 2: Saturated, Unsaturated, or Supersaturated?

These questions test your ability to compare a specific data point to the curve. For example, "If you add 80 grams of NaNO<sub>3</sub> to 100g of water at 30°C, is the solution saturated, unsaturated, or supersaturated?"

**How to solve it:**

1. Find the temperature (30°C) on the x-axis and draw a line up.
2. Find the mass of the solute (80g) on the y-axis and draw a line across.
3. Mark the point where these two lines would meet. Let us call this your "data point."
4. Now, look at where the curve for the substance (NaNO<sub>3</sub>) is at 30°C. Typically, the curve for NaNO<sub>3</sub> at 30°C is around 96 g/100g H<sub>2</sub>O.
5. Compare your data point (80g) to the curve value (96g). Your point (80g) is below the curve (because 80 is less than 96).

**Answer on the worksheet:** Unsaturated. Because you added less solute (80g) than is needed to reach the saturation point (96g), the solution can still dissolve more NaNO<sub>3</sub>.

If the data point was \*on\* the curve, the answer is "saturated." If it was \*above\* the curve, the answer is "supersaturated," which is usually an unstable condition achieved by carefully cooling a hot saturated solution.

## Question Type 3: Determining the Mass of Solute for Crystallization

This is a more advanced question. "A saturated solution of KNO<sub>3</sub> is prepared at 60°C and then cooled to 20°C. How many grams of KNO<sub>3</sub> will crystallize out of the solution?"

**How to solve it:**

1. **Find the solubility at the starting temperature (60°C):** Read the KNO<sub>3</sub> curve at 60°C. This is approximately 108 g/100g H<sub>2</sub>O. So, the hot solution contains 108g of dissolved KNO<sub>3</sub>.
2. **Find the solubility at the final temperature (20°C):** Read the KNO<sub>3</sub> curve at 20°C. This is approximately 32 g/100g H<sub>2</sub>O. At this lower temperature, the water can only keep 32g of KNO<sub>3</sub> dissolved.
3. **Calculate the difference:** Subtract the smaller value from the larger value. 108g (at 60°C) - 32g (at 20°C) = 76g.

**Answer on the worksheet:** 76 grams of KNO<sub>3</sub> will crystallize out of the solution. This mass of solid is "excess" and can no longer be held in the solution, so it forms crystals at the bottom of the container.

## Question Type 4: Comparing Solubility of Different Substances

These questions are designed to test your ability to read multiple curves on a single graph. "At what temperature do KCl and KNO<sub>3</sub> have the same solubility?"

**How to solve it:**

1. Look for the point on the graph where the lines for the two substances (KCl and KNO<sub>3</sub>) intersect or cross each other.
2. Find the temperature on the x-axis at that intersection point.
3. You can also check the y-axis at that point to see what the shared solubility value is.

**Answer on the worksheet:** Approximately 50°C, at a solubility of about 80 g/100g H<sub>2</sub>O.

PRACTICAL EXAMPLES: WORKING THROUGH A SOLUBILITY CURVE WORKSHEET

Let us apply our knowledge to a simulated section from a **solubility curve worksheet with answers**. Imagine a graph with curves for NaCl, KCl, KNO<sub>3</sub>, and Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>. Here are some practice problems and their solutions.

**Problem 1:** How many grams of KCl can be dissolved in 100g of water at 40°C to form a saturated solution?

**Solution:** Find 40°C on the x-axis. Go up to the KCl curve. Trace horizontally to the y-axis. The value is approximately 40g. So, you can dissolve 40g of KCl.

**Problem 2:** A solution is made by adding 110g of KNO<sub>3</sub> to 100g of water at 50°C. Is this solution saturated? Explain your reasoning.

**Solution:** At 50°C, the solubility of KNO<sub>3</sub> is about 85g/100g H<sub>2</sub>O. Since you added 110g, which is more than 85g, the solution is saturated. In fact, 110g is above the curve, so the excess 25g (110-85) would not dissolve and would sit as solid at the bottom. This is a saturated solution with

undissolved solute.

**Problem 3:** Which substance's solubility is least affected by a change in temperature?

**Solution:** Look at the curves. The flattest line indicates the substance whose solubility changes the least with temperature. In most standard graphs, this is sodium chloride (NaCl). Its curve is nearly horizontal. Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> is a trickier one because it often slopes \*downward\*, but its slope is usually steeper than NaCl's flatness. The answer here is NaCl.

**Problem 4:** At 10°C, which solute has the lowest solubility?

**Solution:** Of the common curves, at very low temperatures like 10°C, KNO<sub>3</sub> often has the lowest solubility (around 25g/100g H<sub>2</sub>O), while NaCl has a moderate one (around 35g/100g H<sub>2</sub>O), and Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> is higher. The specific answer depends on the exact graph provided in your **solubility curve worksheet with answers**, but the method is to look at the lowest point on the y-axis for all the curves at the